



Talking to Your Doctor About Plaque Psoriasis

If you experience any of the symptoms of chronic severe plaque psoriasis, make sure you speak to a dermatologist as soon as possible. Remember, only a qualified physician can make a diagnosis and help you make an appropriate treatment choice.

As you speak to your doctor about how plaque psoriasis may be affecting you, use the following questions to guide your discussion. And when the discussion turns to your treatment options, ask your doctor if REMICADE[®] is right for you.

REMICADE[®] is approved for the treatment of adult patients with chronic severe (extensive and/or disabling) plaque psoriasis under the care of a physician who will determine if REMICADE[®] is appropriate considering other available therapies. **Please read the Important Safety Information for REMICADE[®] below.**

Discuss these questions and answers with your dermatologist.

1. What treatments are you taking, or have you taken, in the past?

- Topical ointments/creams
- Phototherapy/PUVA
- Systemics/prescription pills (eg, methotrexate)
- Biologics
- Other

2. How satisfied are you with the results of these treatments?

- Not at all A little A lot

3. How sore, painful, or stinging has your skin felt?

- Not at all A little A lot

4. How much has your skin condition influenced the clothes you wear?

- Not at all A little A lot

5. How strongly do you agree with the following statement?: I am satisfied with my treatment for plaque psoriasis.

- Not at all A little A lot

6. How strongly do you agree with this statement?: I want to try another treatment for my plaque psoriasis.

- Not at all A little A lot

Have some questions of your own?

If you have more questions you'd like to ask, use the space below to write them down. It's always best to prepare your questions before your appointment so you don't forget anything. Let your doctor know at the beginning of the appointment that you have some questions so there will be time to answer all of them.

IMPORTANT SAFETY INFORMATION FOR REMICADE® (infliximab)

Only your doctor can recommend a course of treatment after checking your health condition. REMICADE® (infliximab) can cause serious side effects such as lowering your ability to fight infections. **Some patients, especially those 65 years and older, have had serious infections caused by viruses, fungi or bacteria that have spread throughout the body, including tuberculosis (TB) and histoplasmosis. Some of these infections have been fatal. Your doctor should monitor you closely for signs and symptoms of TB during treatment with REMICADE®.**

Unusual cancers have been reported in children and teenage patients taking TNF-blocker medicines. Hepatosplenic T-cell lymphoma, a rare form of fatal lymphoma, has occurred mostly in teenage or young adult males with Crohn's disease or ulcerative colitis who were taking REMICADE® and azathioprine or 6-mercaptopurine. For children and adults taking TNF blockers, including REMICADE®, the chances of getting lymphoma or other cancers may increase.

You should discuss any concerns about your health and medical care with your doctor.

What should I tell my doctor before I take REMICADE®?

You should let your doctor know if you have or ever had any of the following:

- Tuberculosis (TB) or have been near someone who has TB. Your doctor will check you for TB with a skin test. If you have latent (inactive) TB, you will begin TB treatment before you start REMICADE®.
- Lived in a region where certain fungal infections like histoplasmosis or coccidioidomycosis are common.
- Infections that keep coming back, have diabetes or an immune system problem.
- Any type of cancer or a risk factor for developing cancer, for example, chronic obstructive pulmonary disease (COPD) or had phototherapy for psoriasis.
- Heart failure or any heart condition. Many people with heart failure should not take REMICADE®.
- Hepatitis B virus (HBV) infection or think you may be a carrier of HBV. Your doctor will test you for HBV.
- Nervous system disorders (like multiple sclerosis or Guillain-Barré syndrome).

(Continued...)

IMPORTANT SAFETY INFORMATION FOR REMICADE® (infliximab) (continued)

Also tell your doctor if you:

- Use the medicines Kineret (anakinra), Orencia (abatacept) or Actemra (tocilizumab) or other medicines called biologics used to treat the same problems as REMICADE®.
- Are pregnant, plan to become pregnant, are breast-feeding, or have a baby and were using REMICADE® during your pregnancy. Tell your baby's doctor about your REMICADE® use before the baby receives any vaccine because of an increased risk of infection for up to 6 months after your last dose of REMICADE® you received during your pregnancy.
- Recently received or are scheduled to receive a vaccine. Adults and children taking REMICADE® should not receive live vaccines or treatment with a weakened bacteria (such as BCG for bladder cancer) while taking REMICADE®.

What should I watch for and talk to my doctor about before or while taking REMICADE®?

The following serious (sometimes fatal) side effects have been reported in people taking REMICADE®.

You should tell your doctor right away if you have any of the signs listed below:

- Infections (like TB, blood infections, pneumonia)—fever, tiredness, cough, flu, or warm, red or painful skin or any open sores. REMICADE® can make you more likely to get an infection or make any infection that you have worse.
- Lymphoma, or any other cancers in adults and children.
- Skin cancer—any changes in or growths on your skin.
- Heart failure—new or worsening symptoms, such as shortness of breath, swelling of your ankles or feet, or sudden weight gain.
- Reactivation of HBV—feeling unwell, poor appetite, tiredness, fever, skin rash and/or joint pain.
- Liver injury—jaundice (yellow skin and eyes), dark brown urine, right-sided abdominal pain, fever, or severe tiredness.
- Blood disorders—fever that doesn't go away, bruising, bleeding or severe paleness.
- Nervous system disorders—numbness, weakness, tingling, changes in your vision or seizures.
- Allergic reactions during or after the infusion—hives, difficulty breathing, chest pain, high or low blood pressure, swelling of face and hands, and fever or chills.
- Lupus-like syndrome—chest discomfort or pain that does not go away, shortness of breath, joint pain, rash on the cheeks or arms that gets worse in the sun.
- Psoriasis—new or worsening psoriasis such as red scaly patches or raised bumps on the skin that are filled with pus.

The more common side effects with REMICADE® are respiratory infections (that may include sinus infections and sore throat), headache, rash, coughing and stomach pain.

Click here to read the [Medication Guide](#) for REMICADE® and discuss it with your doctor.
Click here for full [Prescribing Information](#).

You are encouraged to report negative side effects of prescription drugs to the FDA.
Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

026819-141222